



# Pliocene - Quaternary subsidence and exhumation of the Kern arch, southwestern Sierra Nevada, CA, linked to mantle lithosphere removal

Robinson Cecil, Zorka Saleeby, Jason Saleeby, Ken Farley  
California Institute of Technology, Pasadena, CA USA




## 1. Introduction and Motivation

Thermomechanical models of mantle lithosphere removal from beneath the southern Sierra Nevada region, California, predict a complex spatio-temporal pattern of vertical surface displacements. We evaluate these models by using (U-Th)/He thermochronometry, together with other paleothermometry estimates, to investigate such topographic transients. We target strata of the Kern arch, a crescent-shaped uplift located in the SE San Joaquin Basin. Kern arch stratigraphy provides a unique record of subsidence and exhumation in a sensitive region immediately adjacent to the delaminating mantle lithosphere at depth.





Conceptual model of the 3-D delamination of the arclogite root from the Sierra Nevada batholith. A: Early stages of regional east to west delamination. B: hypothetical necking off of a mega-boundin, which promoted ca. 3.5 Ma volcanism. C: initiation of S to N components along southern end of residual root. D: continued N to S delamination progressing to the current state of the Isabella anomaly suspended southeastward into the deeper mantle from the area of residual root attachment under Tulare Basin. Figure from Saleeby et al. 2012.

## 2. Delamination and vertical surface displacements




## 3. The Kern Arch



The Kern arch is a crescent-shaped uplift located along the low western flank of the southern Sierra Nevada. It is adjacent to the Tulare Basin, an area of modern anomalous subsidence centered above the area of residual crustal attachment of the delaminating arclogite at depth. Basinal deposits of the Kern arch region comprise up to 4 km of Eocene - Pleistocene marine and nonmarine strata, which unconformably overlie Early Cretaceous Sierra Nevada basement. Although it has been proposed that the Kern arch is a basement salient which existed since the Paleocene, we argue that it is an entirely recent (Neogene) feature that is associated with Plio-Quaternary removal of mantle lithosphere from beneath the region.

## 6. Forward modeling of detrital apatite (U-Th)/He ages




Well	Elevation (m)	Range of sample collection depths (m)	Range of peak Temps (°C)	Average post-6 Ma sub. (m)	Average post-1 Ma exh. (m)
Fuhrman #1	285	1100 - 1510	70 - 85	1500 <sup>a</sup>	1346 <sup>a</sup>
Richfield SP #15-1	317	1270 - 1540	83 - 88	2425	1732
Dyer Creek #48 <sup>b</sup>	248	988	73	1790	1580
Parsons #1	117	1600 - 2020	84 - 89	1187 <sup>a</sup>	1110 <sup>a</sup>
Smoot #1	239	335 - 354	69 - 71	1655	1828


Subsidence and exhumation estimates tabulated for all wells studied. Estimates from the Smoot-1 well are based on vitrinite reflectance data, which indicated that rocks as shallow as ~ 350 m today were buried to temperatures of 69 - 71 °C.

These estimates are also represented in subsidence curves for the Kern arch shown below. Note that the timing of rapid subsidence and uplift of the arch is coincident with increasing subsidence rates in the Tulare Basin.

## 4. Cryptic subsidence in the Kern Arch



## 5. Thermal history simulations



## 8. References

Gogus and Pysklywec, 2008. Near-surface diagnostics of dripping or delaminating lithosphere. JGR-Solid Earth, v. 113.  
Flowers et al., 2009. Apatite (U-Th)/He thermochronometry using RDAAM. GCA, v. 73, p. 2347-2365.  
Le Pourhiet et al., 2006. Mantle instability beneath the Sierra Nevada mountains in California. EPSL, v. 251, p. 104 - 119.  
Reeg, 2008. Tomographic observations connecting convective downwellings with lithospheric source regions, Sierra Nevada. EOS Trans. AGU, v. 89(53).  
Saleeby et al., 2012. Epeirogenic transients related to mantle lithosphere removal in the southern Sierra Nevada region, California, Part I: Implications of thermo-mechanical modeling, Geosphere.

## 7. Surface transients in the Kern arch - Tulare Basin region

